



Original research

Immediate pain relief effect of low level laser therapy for sports injuries: Randomized, double-blind placebo clinical trial



A. Takenori^{a,*}, M. Ikuhiro^{b,c,d}, U. Shogo^{b,c}, K. Hiroe^e, S. Junji^a, T. Yasutaka^a, K. Hiroya^d, N. Miki^f

^a Faculty of Sports Science, Kyushu Kyoritsu University, Japan

^b Faculty of Physical Education, Osaka University of Health and Sport Sciences, Japan

^c Graduate School of Sport and Exercise Sciences, Osaka University of Health and Sport Sciences, Japan

^d Osaka University of Health and Sport Sciences Clinic, Japan

^e Faculty of Health and Sciences, Tokyo Ariake University of Medical and Health Sciences, Japan

^f Graduate School of Comprehensive Human Sciences, University of Tsukuba, Japan

ARTICLE INFO

Article history:

Received 5 September 2015

Received in revised form 21 January 2016

Accepted 15 March 2016

Available online 24 March 2016

Keywords:

Physical therapies

Athletes

Motion pain

Pain relief rate

ABSTRACT

Objectives: To determine the immediate pain relief effect of low-level laser therapy on sports injuries in athletes and degree of pain relief by the therapy.

Design: Double-blind, randomized, comparative clinical study.

Methods: Participants were 32 college athletes with motion pain at a defined site. Participants were randomized into two groups in which the tested or placebo laser therapy was administered to determine pain intensity from painful action before and after laser irradiation, using the Modified Numerical Rating Scale. The post-therapeutic Modified Numerical Rating Scale score was subtracted from the pre-therapeutic Modified Numerical Rating Scale score to determine pain intensity difference, and the rate of pain intensity difference to pre-therapeutic Modified Numerical Rating Scale was calculated as pain relief rate.

Results: Low-level laser therapy was effective in 75% of the laser group, whereas it was not effective in the placebo group, indicating a significant difference in favor of the laser group ($p < 0.001$). Pain relief rate was significantly higher in the laser group than in the placebo group (36.94% vs. 8.20%, respectively, $p < 0.001$), with the difference in pain relief rate being 28.74%.

Conclusions: Low-level laser therapy provided an immediate pain relief effect, reducing pain by 28.74%. It was effective for pain relief in 75% of participants.

© 2016 Sports Medicine Australia. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Sports injuries constitute a serious problem for many athletes and others who participate in sports because they cause pain and dysfunction, resulting in the inability to continue sports activities. Various physical therapies, including electrotherapy, thermotherapy, cryotherapy, and phototherapy, have been used to alleviate symptoms of sports injuries such as pain.^{1–3} Low-level laser therapy (LLLT) has been clinically introduced as one of such physical therapies.

LLLT has been examined in clinical research and reported to be effective for its long-term effect on many diseases in the general adult population.^{4–6} Bjordal et al.⁷ reported that a single session of LLLT relieved tenderness at the affected site in patients with Achilles tendinitis, and they demonstrated both immediate and long-term effects on injuries in the general adult population.

Studies on the effect of LLLT on sports injuries in athletes are limited to the reports of its effect on sprained ankles⁸ and Achilles tendinopathy.⁹ In both of these studies, the effect of LLLT was the same as was observed in the general adult population. In the study by Stergioulas⁸ in patients with sprained ankles, LLLT was given twice daily significantly alleviated edema at 24–72 h as compared with placebo therapy. In another study by Stergioulas et al.⁹ in recreational athletes with Achilles tendinopathy, the combination of eccentric exercise and LLLT for 4–12 weeks alleviated

* Corresponding author.

E-mail address: awtn9831@gmail.com (A. Takenori).

motion pain as compared with placebo therapy. Thus, LLLT has been demonstrated to alleviate edema and pain associated with sports injuries in a few days to weeks. However, these studies did not provide any data on the immediate pain relief effect of LLLT on sports injuries in athletes.

Because athletes with sports injuries need earlier functional recovery compared to members of the general population, the immediate effect of LLLT is important. Therefore, this study was designed to evaluate whether LLLT provides an immediate pain relief effect on sports injuries in athletes and to determine the extent of pain relief by LLLT.

2. Materials and methods

A double-blind, randomized, placebo-controlled, parallel-group comparison study was performed. Participants were randomly assigned to the laser or placebo group.

Forty-seven college athletes met the following inclusion criteria: participation in intercollegiate to athletic activities 5 days/week or more; treatment at Osaka University of Health and Sport Sciences Clinic between July 1, 2013, and January 31, 2015 for sports injury; and diagnosis by an orthopedist with an orthopedic sports injury for which LLLT was indicated. LLLT was indicated for sport injuries if the following criteria were met: the injuries were painful in motion; the painful area was defined; LLLT was not contraindicated; and the injuries were not associated with any neurological findings.

Exclusion criteria were the inability to define the painful area and absence of definite motion pain. Of the 47 patients enrolled in the study, 32 with a definite painful area and motion pain were included as participants.

The 32 participants were randomly assigned to one of two groups, in which either LLLT or placebo laser therapy was administered (the laser and placebo groups, respectively) according to an assignment table prepared by the coordinator using computer-generated random numbers. The following additional data were collected for each participant: name and site of injury, and period from injury to therapy (number of days after injury). The laser group includes 9 patients with ankle sprain, 1 patient with navicular stress fracture, 1 patient with plantar fasciitis, 1 patient with patella tendinitis, 1 patient with spondylolysis, 1 patient with shoulder arthroscopic surgery, 1 patient with triangular fibrocartilage complex injury, and 1 patient with proximal thumb avulsion fracture. The placebo group includes 5 patients with ankle sprain, 2 patients with meniscal injuries, 2 patients with elbow medial collateral ligament sprain, 2 patients with Achilles tendinitis, 1 patient with low back pain, 1 patient with lumbar facet arthritis, 1 patient with infraspinatus muscle injury, 1 patient with deltoid muscle injury, and 1 patient with shoulder periartthritis.

The sample size of the study was calculated at 15 per group using a statistical power of 0.9, intergroup difference of 30, standard deviation of 25, and significant level of 5% with reference to the results of Malliaropoulos et al.¹⁰ EZR statistical software¹¹ (Saitama Medical Center, Jichi Medical University, Japan, <http://www.jichi.ac.jp/saitama-sct/SaitamaHP.files/statmedEN.html>) was used for calculating the sample size.

This study was performed with the approval of the Research Ethics Committee of Osaka University of Health and Sport Sciences (Approval No. 12-29). Participants who received oral and written explanation of the study and provided written consent to participate were included in the study. LLLT was performed as one of the therapeutic measures after the experiment was completed. Data were collected in the physiotherapy room of Osaka University of Health and Sport Sciences Clinic. None of the participants prematurely discontinued the experiment.

Table 1
Laser parameters.

Wavelength	810 nm (GaAlAs laser)
Frequency	Continuous output
Optical output	180 mW
Spot diameter	0.0007 cm, 0.0005 cm
Spot size	0.0035 cm ²
Power density	51.4 W/cm ²
Energy	5.4 J at each spot
Energy density	1542.85 J/cm ² at each spot
Treatment time	30 s × 20 times (total 10 min)
Irradiation site	1 cm ²
Application mode	Probe held stationary in skin contact with a 90° angle and slight pressure

Participants in the laser group received LLLT from laser therapy equipment (Softlaser JQ-W1, Minato Medical Science Co., Ltd, Japan) with an output of 180 mW, irradiation time of 30 s, and total irradiation time of 10 min (Table 1). Participants in the placebo group received placebo therapy from a placebo device (detuned laser) with an output of 0 mW, irradiation time of 30 s, and total irradiation time of 10 min. The Softlaser used for this study was contact-type laser therapy equipment with an irradiation area of 0.0035 cm². The most painful area during a painful motion was selected as the irradiation site. In order to find the most painful area, participants were asked to explain the most painful motion during their daily or athletic activities. Then participants were asked to identify the most painful area by their index finger during the movement.

Because the study was double-blinded, the measurer and participants were blinded as to whether they used the actual or placebo laser equipment. The output, irradiation time, and total irradiation time of the laser therapy equipment were setup by the coordinator before each participant entered the physiotherapy room. The measurer left the physiotherapy room before the coordinator setup the laser therapy equipment and was call back after the setup. Each participant operated the laser equipment independently after receiving instructions on how to use it by the coordinator. Irradiation site was kept within 1 cm². Therefore a laser probe was applied on the most painful area within a 1 cm² area for 30 s each time for 20 times, and total irradiation time of 10 min. To ensure participant safety and eliminate participant bias, participants were instructed not to look at the laser light during laser irradiation. The measurer observed the therapy procedure and measured pain intensity of the painful motion before and after laser irradiation in both groups. Pain during the painful motion was measured using the Modified Numerical Rating Scale (MNRS), which is a 10-cm scale from 0 to 10 at 1-cm intervals in millimeters, with 0 representing no pain and 10 representing the worst pain.

Injury sites were classified into upper limbs, lower limbs, and body trunk. The MNRS score after the therapy (post-MNRS) was subtracted from that before the therapy (pre-MNRS) to determine the pain intensity difference (PID). The rate of PID relative to the pre-MNRS was calculated as the pain relief rate (PRR).

$$\text{PID} = \text{Pre-MNRS} - \text{Post-MNRS}$$

$$\text{PRR} = \text{PID} / \text{Pre-MNRS} \times 100$$

Statistical analysis was performed using SPSS 21.0J for Windows (IBM Corporation, Armonk, NY, USA) and EZR. The mean number of days after injury, mean PRR, and the 95% confidence interval (95% CI) were calculated for each group.

The difference in injury sites between the groups was tested using Fisher's exact test, with a significance level of 5%. The difference in the number of days after injury, pre-MNRS, or PRR between the groups was tested using an unpaired *t*-test with a significance level of 5%. When a significant difference in the PRR was observed between the groups, PRR was classified into poor, fair, good,

Table 2
Participants characteristics with statistical comparison.

	Laser (n = 16)	Placebo (n = 16)	Difference	p value
Age (years), mean ± SD	20.25 ± 1.18	20.88 ± 2.25		0.335 [*]
Sex (n), male/female	7/9	7/9		1.00 [*]
Height (cm), mean ± SD	164.86 ± 8.97	170.56 ± 9.12		0.085 [*]
Weight (kg), mean ± SD	60.53 ± 11.66	65.68 ± 10.80		0.205 [*]
Pain (MNRS), mean (95%CI)				
Pre	5.29 (3.98–6.61)	5.88 (5.09–6.66)		0.426 [*]
Post	3.26 (2.35–4.18)	5.32 (4.64–6.00)	2.06 (0.97–3.15)	<0.001 [†] (t = 3.850)
PRR (%) mean (95%CI)	36.94 (25.81–48.07)	8.20 (2.43–13.98)	28.74 (16.72–40.75)	<0.001 [†] (t = 4.886)
Injury sites (n)				0.556 [†]
Upper	3	5		
Lower	12	9		
Trunk	1	2		
After injury (day)	37.69 ± 43.36	32.38 ± 35.21		0.706 [*]

MNRS: the Modified Numerical Rating Scale; PRR: pain relief rate.

^{*} Student's *t*-test.

[†] Fisher's exact test.

excellent based on the mean value of the difference and 95% CI. A PRR value equal to or above the mean value of the difference in PRR between the groups was considered to indicate that LLLT was effective, and a PRR value below the mean value was considered to indicate that LLLT was ineffective. The difference in the rate of participants in whom LLLT was effective or ineffective was tested with the χ^2 test with a significance level of 5%.

3. Results

Their characteristics and comparison are shown in Table 2. No significant difference in injury sites ($p = 0.556$, Table 2) or number of days after injury ($p = 0.706$, $t = 0.380$, Table 2) was observed between the groups. The rate of participants in whom LLLT was effective was 75% in the laser group and 0% in the placebo group, and the rate of participants in whom LLLT was ineffective was 25% in the laser group and 100% in the placebo group, with a significant difference between the groups ($p < 0.001$, $\chi^2 = 19.20$, Fig. 1). The PRR was significantly higher in the laser group than in the placebo group (36.94% [95% CI: 25.81–48.07] vs. 8.20% [95% CI: 2.43–13.98]; $p < 0.001$, $t = 4.886$, Table 2), with an intergroup difference of 28.74% (95% CI: 16.72–40.75, Table 2).

4. Discussion

Although an earlier study⁹ showed that long-term LLLT had a pain relief effect on Achilles tendinopathy in sports injuries, no studies have specifically examined the immediate pain relief effect and PRR of LLLT in injured athletes who need early return to play. Therefore, to examine the immediate pain relief effect of LLLT on sports injuries, this study was performed as a double-blind, randomized, placebo-controlled clinical trial to compare the pain relief effect of one session of LLLT in the laser group with that of placebo therapy in the placebo group. The results showed a significantly higher rate of pain relief effect among participants in the laser group ($p < 0.001$). Low-level laser has been reported to provide various effects, including inhibition of nerve excitement,^{12–14} anti-inflammation,^{7,15,16} and tissue repair.^{17–19} The immediate pain relief effect that we observed is more likely related to inhibition of nerve excitement than to chronic effects such as anti-inflammation and tissue repair. In 25% of the laser group in whom LLLT was not effective, no specific characteristics were observed and no specific causal factors could be identified. This is consistent with the results of Malliaropoulos et al.,¹⁰ who reported that 4-weeks LLLT was not effective in 12.5% of patients with meniscal injuries and that no




Fig. 1. PRR classification and comparison of ineffective with effective. PRR: pain relief rate (%). χ^2 test with a significance level of 5%. PRR was classified into poor, fair, good, excellent based on the mean value of the difference and 95% CI. A PRR value equal to or above the mean value of the difference in PRR between the groups was considered to indicate that LLLT was effective, and a PRR value below the mean value was considered to indicate that LLLT was ineffective.

specific causal factors could be determined. In addition to the pain relief effect of LLLT, we examined the PRR to determine the degree of pain relief. The PRR was significantly higher in the laser group than in the placebo group (36.94% vs. 8.2%, respectively, $p < 0.001$), and LLLT immediately relieved motion pain by 28.74% (Table 2). Malliaropoulos et al.¹⁰ reported that 4-weeks LLLT and placebo laser relieved pain by approximately 65% and 22%, respectively, in a randomized controlled trial in patients with meniscal injuries. Bjordal et al.⁷ investigated the change in tenderness at the affected site after a single session of LLLT and reported that LLLT significantly relieved tenderness as compared with placebo laser therapy. The results of the present study support the earlier report of Bjordal et al.⁷ in terms of the comparable immediate pain relief effect of LLLT although the pain relief effect was lower than that observed with 4-weeks LLLT in the study by the Malliaropoulos et al.¹⁰ In the present study, no significant difference was noted between the two groups for injury sites, number of days after injury, or pre-MNRS, indicating that the observed pain relief effect was attributable to low-level laser irradiation and that the effect of injury sites, number of days after injury, and pre-treatment pain intensity was limited. Moreover, LLLT relieved the pain associated with the sports injuries by 36.94%. Based on the difference from placebo therapy, the rate of pain relief from low-level laser irradiation was 28.74%, and immediate pain relief occurred in 75% of the participants.

Bjordal et al.⁷ reported that a single session of LLLT reduced the accumulation of prostaglandin E2, an inflammatory substance, at 105 min. Thus, LLLT appears to have not only an immediate pain relief effect, but also late anti-inflammatory and tissue-repair promoting effects, and may be useful in combination with active exercise therapy. In fact, Stergioulas²⁰ reported that the combination of plyometric exercise and LLLT relieved pain more than the combination of the exercise therapy and placebo laser in patients with humeral lateral epicondylitis. Furthermore, Stergioulas et al.⁹ reported that the combination of eccentric exercise and LLLT resulted in better relief of motion pain than the combination of the exercise therapy and placebo laser in recreational athletes with Achilles tendinopathy. Thus, the present study provides important data to demonstrate the uses (e.g., immediate pain relief and promotion of exercise therapy) and efficacy of LLLT.

The present study was limited by its failure to cover all sports injuries. LLLT was not shown to be effective for all sports injuries.²¹

5. Conclusions

To examine the immediate pain relief effect of LLLT on sports injuries, the present study was performed as a randomized, double-blind, placebo-controlled, clinical study comparing the pain relief effect of a single session of LLLT with that of placebo laser therapy. The results revealed that LLLT relieved pain associated with sports injuries by 36.94%. Based on the difference from the placebo therapy, the rate of pain relief from low-level laser irradiation was 28.74%, and significant immediate pain relief occurred in 75% of the participants. These results serve as important data to demonstrate the usefulness of LLLT in combination with active exercise therapy.

Practical implications

- Low-level laser therapy has immediate effect of pain relief for motion pain in sports injuries.
- Low-level laser therapy should be applied to a small treatment area for pain relief.

- Low-level laser therapy should not apply to the patients with inability to define the painful area and absence of definite motion pain.

Acknowledgements

The authors received no assistance, including research grants, except for the placebo laser device lent from Minato Medical Science Co., Ltd. The authors thank the athletes who participated as subjects in this study.

References

1. Stasinopoulos D. Treatment of spondylolysis with external electrical stimulation in young athletes: a critical literature review. *Br J Sports Med* 2004; 38(3):352–354.
2. Petersen W, Rembitzki IV, Koppenburg AG et al. Treatment of acute ankle ligament injuries: a systematic review. *Arch Orthop Trauma Surg* 2013; 133(8):1129–1141.
3. van den Bekerom MP, Kerkhoffs GM, McCollum GA et al. Management of acute lateral ankle ligament injury in the athlete. *Knee Surg Sports Traumatol Arthrosc* 2013; 21(6):1390–1395.
4. Bjordal JM, Lopes-Martins RA, Joensen J et al. A systematic review with procedural assessments and meta-analysis of low level laser therapy in lateral elbow tendinopathy (tennis elbow). *BMC Musculoskelet Disord* 2008; 9: 75.
5. Chow RT, Johnson MI, Lopes-Martins RA et al. Efficacy of low-level laser therapy in the management of neck pain: a systematic review and meta-analysis of randomised placebo or active-treatment controlled trials. *Lancet* 2009; 374:1897–1908.
6. Chang WD, Lee CL, Lin HY et al. A meta-analysis of clinical effects of low-level laser therapy on temporomandibular joint pain. *J Phys Ther Sci* 2014; 26(8):1297–1300.
7. Bjordal JM, Lopes-Martins RA, Iversen VV. A randomised, placebo controlled trial of low level laser therapy for activated Achilles tendinitis with microdialysis measurement of peritendinous prostaglandin E2 concentrations. *Br J Sports Med* 2006; 40(1):76–80.
8. Stergioulas A. Low-level laser treatment can reduce edema in second degree ankle sprains. *J Clin Laser Med Surg* 2004; 22(2):125–128.
9. Stergioulas A, Stergioula M, Aarskog R et al. Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. *Am J Sports Med* 2008; 36(5):881–887.
10. Malliaropoulos N, Kiritsi O, Tsitas K et al. Low-level laser therapy in meniscal pathology: a double-blinded placebo-controlled trial. *Lasers Med Sci* 2013; 28(4):1183–1188.
11. Kanda Y. Investigation of the freely available easy-to-use software 'EZ' for medical statistics. *Bone Marrow Transplant* 2013; 48(3):452–458.
12. Tsuchiya K, Kawatani M, Takeshige C et al. Diode laser irradiation selectively diminishes slow component of axonal volleys to dorsal roots from the saphenous nerve in the rat. *Neurosci Lett* 1993; 161(1):65–68.
13. Tsuchiya K, Kawatani M, Takeshige C et al. Laser irradiation abates neuronal responses to nociceptive stimulation of rat-paw skin. *Brain Res Bull* 1994; 34(4):369–374.
14. Kasai S, Kono T, Yamamoto Y et al. Effect of low-power laser irradiation on impulse conduction in anesthetized rabbits. *J Clin Laser Med Surg* 1996; 14(3):107–109.
15. Hegedus B, Viharos L, Gervain M et al. The effect of low-level laser in knee osteoarthritis: a double-blind, randomized, placebo-controlled trial. *Photomed Laser Surg* 2009; 27(4):577–584.
16. Simunovic-Soskic M, Pezelj-Ribaric S, Brumini G et al. Salivary levels of TNF-alpha and IL-6 in patients with denture stomatitis before and after laser phototherapy. *Photomed Laser Surg* 2010; 28(2):189–193.
17. Fung DT, Ng GY, Leung MC et al. Therapeutic low energy laser improves the mechanical strength of repairing medial collateral ligament. *Lasers Surg Med* 2002; 31(2):91–96.
18. Enwemeka CS, Parker JC, Dowdy DS et al. The efficacy of low-power lasers in tissue repair and pain control: a meta-analysis study. *Photomed Laser Surg* 2004; 22(4):323–329.
19. Woodruff LD, Bounkeo JM, Brannon WM et al. The efficacy of laser therapy in wound repair: a meta-analysis of the literature. *Photomed Laser Surg* 2004; 22(3):241–247.
20. Stergioulas A. Effects of low-level laser and plyometric exercises in the treatment of lateral epicondylitis. *Photomed Laser Surg* 2007; 25(3):205–213.
21. Winters M, Eskes M, Weir A et al. Treatment of medial tibial stress syndrome: a systematic review. *Sports Med* 2013; 43(12):1315–1333.